

Contents

Introduction	
Management and Ownership Structure	2
Management Objectives	
Stakeholders	6
Regulations	8
Environmental Policy and Practices	11
Estate Description	12
Forest Description	15
High Conservation Value Areas	17
Establishment and Silviculture	18
Inventory, Mapping and Forest Records	20
Harvesting	22
Protection and Maintenance	23
Monitoring	26
Planning	27
Appendices	28



Introduction

Wood has many different uses, from firewood to construction timber, to newspapers and tissue paper, from furniture to clothing. Wood fibre is extremely versatile and is produced using carbon dioxide and the energy of the sun. Wood from well managed forests is a renewable resource which can meet the economic, social and cultural needs of our society without compromising the environment.

Wairarapa Estate Limited's objective is to grow wood for further processing in New Zealand or overseas and obtain an economic return on investment. Wairarapa Estate seeks to achieve this through the growing of suitable species with wood characteristics which meet the demand of the market. The trees are established, protected and tended as required to meet those demands.

Principles and Criteria

The estate has achieved Forest Stewardship Council® (FSC®) Certification, Licence Code FSC® C131650.

The Forest Owners are committed to the FSC Principles, Criteria and Standards of Good Forest Management. These standards include ecological, social and economic parameters.

The Manager will be able to market the forest log products as certified wood for input into domestic processor markets and international log markets where some are seeking certified wood inputs for manufacture of certifiable end-use products to international consumer markets.

Maintenance of the Certified status is managed through the Manager's Integrated Management System (IMS) and associated documents, IT-based support tools, and by a process of internal and independent external audit undertaken at least annually.

About the Forest Management Plan

This **Forest Management Plan ("the Plan")** is a broad document defining the management structures and vehicles in place to manage an aggregation of individual forests subject to different ownership structures or relationships. Each operational forest, or group of forests, has its own detailed management plan which describes basic but specific management details.

This document provides a summary of the Forest Management Plan for the forests and contains:

- Ownership, management and management planning structure
- Management objectives
- A description of the land and forest resources
- Environmental safeguards
- Identification and protection of rare, threatened and endangered species
- Management regimes and harvest planning
- Forest health
- Management of HCV areas and reserves
- Maps showing plantation area, legal boundaries and protected areas
- Provisions for monitoring and protection



Management and Ownership Structure

The estate is owned by Wairarapa Estate Limited, which is wholly owned by ANZFOF2 NZ Pty Ltd, situated at Level 23, 141 Walker Street, North Sydney, NSW 2060, Australia.

Forest Enterprises Growth Limited has been appointed as Property Manager ("the Manager") by Wairarapa Estate. The estate is a mixture of freehold land and Forestry Rights.

Wairarapa Estate will protect any resource and tenure rights of Tangata Whenua.

As Property Manager, Forest Enterprises will manage day-to-day operational activities and in general manage the estate to standards required to maintain FSC certification.

Forest Enterprises operates out of its head office in Masterton. Forest Management Limited (FML) manages harvest operations in Wairarapa Estate forests in the Manawatu/Wanganui, Hawkes Bay and Tararua regions.

The people responsible for enacting this Plan are:

- Bert Hughes RMNZIF CMInstD CEO and Forestry Director, Forest Enterprises
- Malte Coulmann RMNZIF Forest Estate Manager, Forest Enterprises
- Chris Hawthorn Forest Operations Manager, Forest Enterprises
- Tom Otterson Harvest Manager, Forest Enterprises
- Hannah Harvey Environmental & Risk Co-ordinator, Forest Enterprises
- Stu Harvey GIS Manager, Forest Enterprises
- Bryn Townend GIS Analyst, Forest Enterprises
- John (JJ) Rasmussen Operations Manager, FML
- Gary Connolly Operations Manager, FML
- Pete Newland Harvest/Training Manager, FML



Management Objectives

Economic and Social

The forests are managed to provide environmental benefits, including:

- Enhanced water quality
- Soil stabilisation and conservation
- A buffer against flooding during storms
- Shading waterways for aquatic life
- Enhance wildlife and plant habitat leading to increased biodiversity
- A reduction in greenhouse gases
- Providing economic and social benefits to the community

The forests are all managed to the Manager's Environmental Standards.

Management

The Manager is committed to ensuring the management of the forest is sustainable, achieves economic sustainability and provides the best possible returns for the forest owner. In addition, the Manager will ensure all forests retain the capacity to do the above while meeting a range of environmental, social and cultural outcomes.

The estate will be managed to:

- Ensure the estate is a renewable and sustainable resource
- Grow trees and produce logs for wood products in New Zealand and overseas
- Ensure productivity of the land does not decline
- Ensure environmental and social values are identified and maintained, and undertake operations to minimise impacts on the environment and the community
- Ensure historic sites are identified and protected
- Harvest trees as close as possible to their economic optimum age and achieve the best possible financial returns to the Forest Owners
- Replant following harvesting where land tenure allows
- Meet all statutory requirements for example, New Zealand's Resource Management Act 1991, the National Environmental Standards for Plantation Forestry 2017, the New Zealand Forest Accord 1991, the Principles for Commercial Plantation Forest Management in New Zealand 1995, and comply with forest industry best practice
- A safe and healthy workplace free of workplace injuries
- Act as a good corporate citizen and neighbour
- Ensure all forest management practices are consistent with the principles of the Forest Stewardship Council
- Identify and protect areas of significant ecological and scientific value within the forests and put in places processes to protect and enhance identified values
- Ensure forest sustainably and minimise adverse effects of forest operations on soil and water values
- Minimise impact of operations on archaeological and cultural sites and ensure compliance with the Heritage New Zealand Pouhere Taonga Act 2014
- Minimise impact of operations on amenity values (visual, noise and air effects) and neighbouring properties
- Use chemicals responsibly and seek to minimise the use of chemicals in operations as far as practical



- Capture and learn from environmental incidents through incident reporting, investigation and sharing of learnings
- Ensure staff and contractors receive appropriate training to comply with the law and the requirements of the company Integrated Management System
- Monitor, research and seek new ways to minimise impacts of forestry operations on the environment, and maximise environmental benefits of forests
- Recognise the recreational value of the forest estate to local communities and the general public, and proactively manage public access taking into account safety of people, environmental considerations and forestry operations
- Identify areas within our estate which meet the FSC definition of High Conservation Value Areas and manage these in accordance with FSC requirements
- Ensure there is no conflict with written leases and right of ways as per the individual agreements

Wairarapa Estate and the Manager are committed to ensuring the management of the forest is sustainable from an environmental, social, cultural and economic perspective. These perspectives underpin the FSC management culture.

1. Environmental Perspective

Includes steps to identify rare, threatened and endangered species where such presence is a possibility, protection of reserve areas, waterways and the control of pests and weeds.

2. Social Perspective

Includes ensuring contractors and their workers adhere to health and safety standards, consultation with neighbours and stakeholders in respect of operations on the forest occur. All staff have the right to be a member of a union if they wish.

3. Cultural Perspective

Includes consultation with the appropriate iwi to ensure culturally significant resources, land, historic and archaeological sites are identified and appropriately managed.

Wairarapa Estate will be managed to protect any resource and tenure rights of Tangata Whenua. Tangata Whenua are being identified for each area. However, at time of writing no culturally important areas have been identified.

4. Economic Perspective

Refers to the selection of a species and ensuring management and harvesting regimes provide a reasonable return on investment while minimising the risks of investment.

External Agreements

Through its membership of the New Zealand Forest Owners Association, the Manager is bound by the requirements of the New Zealand Forest Accord 1991 and the Principles for Commercial Plantation Forest Management in New Zealand 1995.

The Forest Accord protects remaining indigenous forest remnants within the plantation forest which meet minimum size and quality criteria from clearance and conversion to plantation forest. All New Zealand Forest Accord vegetation within Wairarapa Estate is identified in a Geographic Information System (GIS) and is protected.

The Principles for Commercial Plantation Forest Management in New Zealand are complementary to the New Zealand Forest Accord. They cover a range of broader principles to promote environmental excellence in plantation forest management, and the protection, preservation and sustainable management of native forests.

The Manager encourages all staff to join the New Zealand Institute of Forestry which requires an annual agreement to maintain a code of conduct and provides professional development opportunities for members.



Implementation

The forest management objectives described above are implemented by the Manager. The Manager applies best forest management practice within a quality management framework to plan for and deliver the required forest management objectives.

The Quality Management Framework includes:

- A forest management system (Tigermoth and GeoMaster) to ensure the forest management planning is up to date and operations are scheduled and undertaken according to the plan
- Environmental Standards to ensure operations follow the Standards, and ensuring high standards of environmental management is integrated into all areas of forest and operational planning and management
- The Health and Safety Management System (HSMS) to ensure all operations are managed safely with the goal of zero serious harm
- FSC Certification to ensure management principles and practice adhere to internationally recognised and adopted standards for forest management



Stakeholders

Community relations are an important focus for the Manager who is committed to being ethically and socially responsible, while meeting its business needs. The Manager makes every attempt to operate with the communities which neighbour Wairarapa Estate forest properties.

The Manager strives to actively engage with stakeholders in the many communities in which we operate, and particularly those directly or indirectly affected by our operations. Prior to commencing harvesting in a new area, the Manager engages with representatives of the local community to keep them informed of plans and develop mitigation strategies for identified concerns. Typically, this includes forest neighbours, residents of any rural access roads affected by logging traffic, and Tangata Whenua.

Social Impact Assessment

The Manager undertakes a social impact assessment annually. This assessment is available to public on request.

The Manager has developed a Social Impact Assessment (SIA) procedure to identify and manage decisions which may have significant impact on the local community. Key staff receive training in SIA techniques.

Key results from the SIA are:

(A) Regional Development

The Manager will remain alert to any adverse impacts from its operations and deal with them accordingly, either individually or as part of an industry grouping. The major impacts are mostly felt by neighbours. The Manager has a list of neighbours and refreshes the list before the start of the Fire Season each year.

Impacts on neighbours are measured by records of compliments and complaints maintained in the Manager's office, records of cooperation on boundary spraying, summer grazing run-off etc.

(B) Optimal Use

Good environmental stewardship means the existing indigenous vegetation enclaves have been assessed, ranked and are protected. Additional areas such as riverine gravels and limestone outcrops are also protected for their environmental values.

(C) Illegal Activities

No illegal logging takes place.

The neighbours referred to in (A) above maintain a good watch over Wairarapa Estate forestlands, mainly to protect their own stock from poachers who may hunt in Wairarapa **Estate's** blocks. The police are also available to assist if required. The Manager's employees have cameras within the forest to monitor illegal

(D) Skills Development

Training plans for staff and contractors will be established each year.

Contractors selected for specific tasks will be selected on the basis of specific criteria including proven skills for the task, training achievements and experience.

The Manager will maintain and be part of industry initiatives to develop a motivated, drug-free and fully skilled workforce for the industry's needs in the near future.

(E) Health & Safety

The Manager has a documented Health & Safety Management System (HSMS).

A formal induction to each forest block and work site is a prerequisite of each operation.

The Manager maintains an Integrated Management System (IMS) incident database and compares itself against the overall industry Loss Time Injury Rate (LTIR) figures. Near misses are reported, as are incidents. Significant near misses will be investigated by the contractor and company supervisor in the spirit of a "no blame" culture. Records are kept and analysed for trends.

Reports on accidents and investigations from other companies are circulated to crew. Ensuring public safety is the rationale behind strict entry controls on Wairarapa Estate forests. There are no staff available to patrol on a regular basis and so free public entry is not an option.



(F) Worker Rights

Crew members have the opportunity to belong to a union.

The Manager ensures any principal contractors used on the estate comply with legislation for minimum wage rates, holidays, superannuation and sick leave. The Manager does not discourage staff or contractors from joining unions should they wish to do so.

(H) Tangata Whenua

Wairarapa Estate will protect any resource and tenure rights of Tangata Whenua.

(I) Neighbours

All neighbours are rural dwellers and are involved in a range of rural economic activities such as farming, tourism, horticulture, agriculture or other small businesses. Corporate neighbours are usually other forest owners.

Neighbour contact details are maintained through databases. Some or all of these parties will be consulted when operations are proposed in forest areas adjacent to their boundaries.

(J) Research Organisations

GNS (Geological and Nuclear Sciences) are very interested in the Saline Springs in Glenburn forest. The manager will ensure they have access to these springs at all times.



Regulations

In order to minimise the risk to forest owners, managers and contractors, it is important relevant legislation and agreements are identified and appropriate measures put in place to ensure breaches of legislation are avoided.

The Manager is kept current on changes in legislation by receiving communications issued by the New Zealand Law Society, reviewing the regular legal updates provided by its external legal advisors, and those published in the New Zealand Institute of Forestry newsletter, as well as reviewing any relevant updates sent out by the New Zealand Forest Owners Association.

All New Zealand legislation is available at http://legislation.govt.nz. Important legislation is in hardcopy within the Managers office and this is reviewed and updated annually as required.

The following legislation and agreements summarise key regulatory and voluntary controls which currently apply to forest operations in the forest.

National Environmental Standards for Plantation Forestry

The estate is subject to the provisions of the National Environmental Standards for Plantation Forestry (NES-PF). The NES-PF is a resource management system under the RMA that provides a nationally consistent set of standards to manage the environmental effects of plantation forestry activities.

Resource Management Act

The estate is subject to the provisions of the Resource Management Act 1991 (RMA). The RMA is a resource management system which promotes the sustainable management of natural and physical resources and is now the principal statute for the management of land, water, soil and other resources in New Zealand.

Under the RMA, each council has its own plans and rules which must be adhered to.

District Councils look after land management issues such as land use, landscapes and biodiversity. Regional Councils deal with soil conservation, water quality issues, discharges to the air, water and land and the coastal marine environment.

Regional Council	District Council
	Hastings
Hawkes Bay	Wairoa
	Central Hawkes Bay
Horizons	Manawatu
Horizons	Tararua
	Masterton
Greater Wellington	Carterton
	South Wairarapa

Heritage New Zealand Act

Under the Heritage New Zealand Pouhere Taonga Act 2014, it is the landowner's responsibility to identify any historic sites on their land prior to undertaking any work which may disturb or destroy such sites. Where such circumstances might exist, an "Authority to Modify or Destroy" will be sought from Historic Places Trust (HPT). Such authorities are similar in function to a resource consent and, if granted, normally come with conditions which must be met.

Records of archaeological and historical places are maintained in the New Zealand Archaeological Association Site Recording Scheme run by the HPT. There is a searchable register maintained online on the HPT website. To search in this register, follow this link:

http://www.historic.org.nz/en/TheRegister/RegisterSearch.aspx

Registered historic sites are also often included in schedules of places and sites of significance in District Plans along with sites of cultural significance.



If a site is found or suspected on any block, the protocols specified in the Manager's Environmental Standards, and any others specifically developed in conjunction with HPT and iwi or other stakeholders must be observed.

Resource Consents

There are currently 28 consented activities which require Resource Consents held for Wairarapa Estate.

Emissions Trade Scheme

Forests in New Zealand are **governed by rules related to New Zealand's commitments to** reduce greenhouse gas emissions.

Any existing forest originally planted prior to 1 January 1990 will be required to cover all their emissions if the forest is deforested. Deforestation occurs if the forest is not replanted, is left to regenerate naturally or, does not achieve the regulated heights and stocking densities as required under the Climate Change Response Act 2002.

All Post-1989 forest in Wairarapa Estate is not registered in **New Zealand's** Emissions Trading Scheme (ETS). Wairarapa Estate does not own Pre-1990 NZUs.

Other Relevant Legislation

- Animal Welfare Act 1999
- Biosecurity Act 1993
- Climate Change Response Act 2002
- Conservation Act 1987
- Fencing Act 1978
- Forests Act 1949
- Fire and Emergency New Zealand Act 2017
- Forests Amendment Act 1993
- Forestry Rights Registration Act 1983
- Hazardous Substances and New Organisms Act 1996
- Health & Safety at Work Act 2015
- Injury Prevention, Rehabilitation and Compensation Act 2001
- New Zealand Forest Accord
- Noxious Plants Act 1978
- Pesticides Act 1999
- Reserves Act 1977
- Soil Conservation and River Control Act 1941
- Trespass Act 1980

Additional relevant legislation is included in the Appendix of this Plan.

Environmental Code of Practice

All operations carried out on the estate must be undertaken to the standards specified in the New Zealand Environmental Code of Practice for Plantation Forestry and the New Zealand Code of Practice for Forest Engineering.

Health and Safety

All operations managed by the Manager are subject to the Manager's Health and Safety Management System (HSMS). This programme includes active accident prevention programmes, training, injury management and drug and alcohol testing.

Health and safety statistics are reported quarterly to Wairarapa Estate Limited.



Responsibilities and Authorities

All staff are responsible for ensuring operations under their immediate control are planned and carried out to meet relevant requirements of any Resource Consent or Permitted Activity Conditions.

Staff are also required to ensure the Manager's crews and/or contractors carrying out these operations are fully aware of these requirements and the steps required to comply.

Any breach of these requirements is deemed a Significant Environmental Event and shall be dealt with as such.

The Environment and Forestry Activities

Forestry activities encompassing silvicultural and harvesting operations can have both beneficial and adverse impacts on the environment depending on the quality of environmental and operational management.

Well managed forests can:

- Enhance water quality
- Stabilise and conserve soil
- Provide a buffer against flood flows during storms
- Shade waterways keeping water cool for enhanced fish and macroinvertebrate life
- Provide habitat for rare, threatened and endangered native species
- Sequester carbon to combat climate change and
- Provide recreational, economic and social benefits to the community

Conversely, poorly managed forestry activities can have harmful impacts. The Manager aims to identify the potential negative impacts and to implement environmental safeguards to prevent or to minimise the negative impact from its operations.



Environmental Policy and Practices

Environmental policy and practices are an integral part of every operation which takes place on the forest. The Manager maintains an Environmental Policy Statement which is signed by the Forestry Director and followed by all staff.

Regular monitoring of key environmental parameters will be undertaken where necessary to ensure the impact on the forest environment from events such as windstorms, flooding and fire, or of agents such as pests, diseases, and weeds are minimised.

The management of the forest recognises the importance of the natural and social environment for the future of its business. The people employed in the forest and processing plants, the neighbouring landowners, the appropriate iwi and the community at large are all recognised as stakeholders.

All activities within the estate are subject to the Manager's Environmental Standards.

Environmental Goals

- 1. Achieve a greater understanding from all persons working within the forest of their environmental responsibilities
- 2. Establish working relationships with all councils
- 3. Establish relationships with iwi, neighbours, and other stakeholders
- 4. Promote and undertake sound environmental stewardship of land and other natural resources on, or adjacent to, this land

Environmental Standards

The Manager's Environmental Standards set out the expectations with regard to managing the environment during forest operations. They are designed to communicate expectations for environmentally sound forestry operations.

Standard Operating Procedures (SOPs) guide all operations and a continuously reviewed and updated.

Assessment of environmental risks due to operations are covered within the SOPs along with other specific forms relating to harvest operations, some of which are managed on BraveGen.

The Environmental Standards and SOPs cover hazardous substances management and clearly outlines the expectations the Manager has of all staff, contractors and suppliers in relation to hazardous substance management.

Hazardous materials which may be used within Wairarapa Estate are:

- Herbicides
- Pesticides
- Fuels
- Oil
- Fire retardants
- Surfactants
- Paint

The Manager is committed to reducing the use of hazardous substances. All aspects of chemical use are reported annually.



Estate Description

Forest Area

Wairarapa Estate is made up of 23 forests with a total Gross area of 10011.20 hectares. The table below shows the breakdown of the estate by forest and type. Wairarapa Estate Ltd. and Southland Estate Ltd. are in the same Estate Fund, therefore are one Forest Management Unit (FMU).

Forest	Gross Area (HA)	Net planted area (ha) All Species	Working Forest Area	FSC RSAA (HA)	RSAA (% of net productive area)
Hapua	743.60	19.41	654.62	39.6	6.04
Waipukurau	77.10	70.79	70.79	2.0	2.80
Craigie Lea	1674.80	1198.82	1198.82	377.5	31.49
Dunolly	602.20	424.8	424.8	75.0	17.66
Driscoll road	155.20	84.86	109.18	14.0	12.81
Erindale	875.80	601.8	657.57	187.2	28.46
Flat Point	154.70	100.21	100.21	47.8	47.72
Glenburn	2031.60	1546.21	1546.21	239.7	15.50
Hawkins	176.10	127.84	147.84	11.5	7.81
Kaiwhata Pines	215.50	145.28	145.28	52.4	36.09
Lands End	117.30	86.27	91.42	4.0	4.41
Ngahape - WE	276.30	137.26	193.47	41.4	21.37
Oldfields	212.80	115.58	177.32	24.7	13.95
Pakowhai	708.00	505.31	572.81	109.0	19.03
Pongaroa	405.80	293.53	293.53	97.3	33.14
Ruakokoputuna	262.90	141.26	141.73	9.2	6.52
Tinui - WE	189.90	142.55	142.55	15.9	11.16
Woodford Green	162.20	67.72	147.61	4.0	2.74
Wai Ngaio	242.70	125.68	128.32	110.9	86.43
Roil	199.20	39.14	129.84	47.0	36.21
Beehive Creek	166.70	106.03	106.03	31.4	29.60
Putorino	188.40	176.25	176.25	2.8	1.57
Riverina	172.40	140.34	156.09	11.2	7.16
WEL Total	10011.20	6396.94	7512.29	1555.47	21%
_	107	100.10	100.15	4= 6	45.00
Fox	127	100.16	100.16	15.6	15.60
Homestead	109.2	85.14	85.14	14.4	16.90
Old Shed	138.9	107.24	107.24	24.4	22.78
Tyneholm	132.1	121.57	121.57	4.1	3.35
Woodslea	122	99.93	99.93	14.6	14.63
SEL Total	629.2	514.04	514.04	73.13	14%
Total by FMU **Cutting/Forestry Right Forest	10640.40	6910.98	8026.33	1628.60	20.29%

^{**}Cutting/Forestry Right Forest

Species

Wairarapa Estate is primarily Pinus radiata, with some minor species planted in patches. All radiata crops are managed either on a pruned, framing or untended regime. It is intended to replant all areas after harvest with improved genetics.



Pinus radiata is the most extensively used species for plantation forestry in New Zealand due to its fast-growing behaviour, durability and versatility. It was introduced to New Zealand in the 1850s, adapted to our climate quickly and by the early 1900's radiata plantations were throughout the country. During the 1950s, genetic improvement commenced and currently continue to produce higher quality wood. Demand for quality wood is always increasing both domestic and internationally.

Age Class


Wairarapa Estate has an uneven age-class spread with the majority of stand aged 21-25 years. The following table shows the breakdown of the estate by age class.

Tree Age (years)	Area (ha)
Fallow	884.5
1-5	780.3
6-10	100.9
11-15	96.5
16-20	228.7
21-25	4291.8
26-30	892.9
30+	43.7

Wairarapa Estate Map

A map of the forests comprising Wairarapa Estate is included on the next page.





Forest Description

Location

Wairarapa Estate currently extends over three regions and eight districts:

Forest Name	Region	District	
Hapua			
Putorino		Hastings	
Roil	Hawkes Bay		
Riverina - Wairoa		Wairoa	
Waipukurau		Central Hawkes Bay	
Beehive Creek	Managari	Manawatu	
Pongaroa	Manawatu-Wanganui	Tararua	
Erindale			
Hawkins			
Ngahape		Masterton	
Oldfields			
Pakowhai			
Woodford Green			
Craigie Lea			
Driscoll road	Wellington		
Dunolly	vveiiiigtori		
Flat Point		Carterton	
Glenburn			
Kaiwhata Pines			
Lands' End			
Ruakokoputuna			
Tinui		South Wairarapa	
Wai Ngaio			

Topography

Hawkes Bay

Hawkes Bay is largely made up of hill country and coastal plains. The Huiarau, Kaweka, and Ruahine Ranges uplift the region (earthquake prone) which cause thin-soiled erosion-prone hills. This land is often marginal and can only be used extensively which makes forestry an appropriate land use.

Manawatu-Wanganui

The Manawatu-Wanganui region has gentle rolling hills to flat coastal plains as well as the steep Tararua and Ruahine Ranges that separate the region from Wellington. The coastal plains are shaped by extensive river systems which makes the region mostly suitable for intensive land use such as dairy farming and cropping, however, steeper land and land needing erosion protection is suitable for plantation forestry.

Wellington

The Wellington Region, like Manawatu-Wanganui, has a variable topography that includes the steep Tararua and Ruahine Ranges that dissipates into rolling hills and plains. The Wairarapa also includes the Aorangi Range which provides regional central uplift.



Geology and Soils

Wairarapa Estate, which lies along the East Coast of New Zealand's North Island, is part of a complex geological accretionary system where faulting and folding of rock is significant due to the subduction of the Pacific Oceanic Plate under the Australian Continental Plate. This subduction has created an accretionary wedge regime along the East Coast of the North Island. The geology of Wairarapa Estate ranges from Carboniferous (300Mya) (Greywacke Tararuas/Ruahine Ranges) to Quaternary (Recent) in age.

The soils of Wairarapa Estate are majority Brown and Pallic soils with some minor Gley, Pumice, Raw, Recent Soils and one area of podzol. Brown Soils are the most extensive soil type for New Zealand, whereas Pallic Soils cover 12% of New Zealand. Brown Soils have relative stable topsoils with well-developed subsoils whereas Pallic Soils have limited permeability and are surface erosion susceptible (ideal for P. Radiata compared to grass species).

For more information on New Zealand Soils - Hewitt, A., & Dymond, J. (2013). Survey of New Zealand soil orders. Ecosystem services in New Zealand: conditions and trends, 1(10), 121-131

Climate

Hawkes Bay

Temperature (C°)	Sunshine	Rainfall	Ground Frost
Mean daily maximum: 19.5 Mean monthly temperature: 14.55 Mean daily minimum: 9.6	Average annual bright sunshine hours: 2238.2	Average monthly rainfall (mm): 74.21 Average monthly wet days (1mm Or More of Rain): 8.15 Mean annual rainfall (mm): 877.35	Average days of frost: 29.65 per annum

Manawatu-Wanganui

Temperature (C°)	Sunshine	Rainfall	Ground Frost
Mean daily maximum: 17.9 Mean monthly temperature: 13.65 Mean daily minimum: 9.3	Average annual bright sunshine hours: 1899.25	Average monthly rainfall (mm): 76.49 Average monthly wet days (1mm Or More of Rain): 9.89 Mean annual rainfall (mm): 918.3	Average days of frost: 23.75 per annum

Wellington

Temperature (C°)	Sunshine	Rainfall	Ground Frost
Temperature (C°) Mean daily maximum: 17 Mean monthly temperature: 12.85 Mean daily minimum: 8.7	Average annual bright sunshine hours: 1982.2	Average monthly rainfall (mm): 88.95 Average monthly wet days (1mm Or More of Rain): 10.51 Mean annual rainfall (mm): 1069.15	Average days of frost: 36 per annum



High Conservation Value Areas

Indigenous biodiversity management within the forests is an essential component of everyday forest management. Although exotic forests can provide a level of biodiversity, High Conservation Value (HCV) areas are usually the source of most indigenous biodiversity. Rare, Threatened and Endangered species (RTE) can also be found associated with exotic forests and require special attention for management.

All reserve areas are managed per Forest Enterprises' High Conservation Value Areas and Reserves Policy.

The Manager contracts Wildlands Ltd to identify HCV areas. As forests are added to the estate, the Manager identifies HCV areas within the new forests. Wildlands Ltd will be contracted every two years to check and monitor Category One and HCVF areas, and all other classified areas every 5 years.

Wairarapa Estate consists of 1550 hectares of Reserve areas. All HCV areas have been mapped and recorded with the Manager's GIS.

Category One Reserve Areas

All these areas have been mapped and identified and recorded on the GIS. The Manager will protect these areas.

Wildlands Ltd will monitor the areas on a two-year cycle for pests, weed and rare and threatened species.

Category Two Reserve Areas

All these areas have been mapped and identified and recorded on the GIS. The Manager will protect these areas; however, the areas can be crossed for operational purposes if this is the best environmental result. Any crossing of the area will require a management plan.

Wildlands Ltd will monitor the areas on a five-year cycle for pests, weed and rare and threatened species.

Category Three and Four Areas

All these areas have been mapped and identified and recorded on the GIS. The Manager will protect these areas; however, the areas can be crossed for operational purposes if this is the best environmental result. Any crossing of the area will require a management plan.

Wildlands Ltd will monitor the areas on a five-year cycle for pests, weed and rare and threatened species.

Rare, Threatened and Endangered Species

All contractors and staff must be trained to identify and given the identification form for (also in Environmental Standards), known rare, threatened and endangered species. If any species are found, the Manager is to be notified immediately and a species sighting form completed.

Protection requirements are assessed at the time of re-establishment where additions to riparian or buffering setbacks are often recommended.

In the case of fauna, records of sighting and locations are stored within GeoMaster and GIS.

Whenever an operation is planned, GeoMaster and GIS must be checked for any record of rare and threatened species within or near the operational area. If there have been observations of any species, all contractors and their staff must be made aware of this, and a management plan for protecting the species will be prepared as part of the Harvesting Prescription. If required, this plan will be prepared in association with local experts.



Establishment and Silviculture

All forest operations are planned to ensure the crop achieves maximum growth and is of high quality.

Typical establishment and silviculture operations within Wairarapa Estate include:

- Land preparation
- Planting
- Weed control
- Pest and disease control
- Fire protection
- Pruning
- Thinning

In addition, the Manager follows a maintenance plan which includes road, track, fence and water way maintenance.

All operations must follow the standards set within the Manager's Environmental Standards.

Establishment

No land under native vegetation will be converted into plantations as per the New Zealand Forest Accord.

The majority of harvested areas will be replanted the winter following the first spring after harvest.

Establishment may include:

- Raking of slash
- Spot mounding
- Aerial desiccation spraying
- Planting of genetically improved seedlings (generally Pinus radiata is planted at 1,000-1,250 stems per hectare)
- Animal pest control
- Fertilising
- Aerial or spot releasing of weed competition

Prior to any forest establishment, a review of the area will be conducted to identify whether there are any risks to rare, threatened or endangered species of flora or fauna. At the same time, consideration will be made of riparian buffer sizes and hard to harvest areas.

All establishment sites will be reviewed to ensure reserves and HCV areas are maintained, that there is a mix of age classes throughout Wairarapa Estate, and correct genotypes are used.

Silviculture

Silviculture is the practice of controlling the growth, composition, health and quality of a forest to meet specific objectives.

There are two main tending regimes within Wairarapa Estate - pruned and framing. Some areas are left untended as a protection crop or for other reasons, however this is not the norm.

Future regimes will depend on Wairarapa Estate and the Manager's assessment of market opportunities, site factors (including slope) and the tree-stocks available.

Thinning

Thinning of stands is undertaken, generally between six to nine years of age, to provide the optimum space for selected crop trees within the stand to grow and maximise their economic return. The aim is to thin out the smaller or poorer formed trees leaving the bigger, better formed trees to grow on. Most thinning operations leave the thinned stems on the forest floor to decompose where production thinning is impractical or uneconomic.



Regimes

Pruned

Year	Operation	Stems per ha	Details
0	Establish	1000-1250	Improved genetics
5-6	Prune 0-3m	375	Minimum green crown must be 3.0m
7-8	Prune 3-6m	375	Minimum green crown must be 3.5m
7-8	Thin to waste	350-375	Thin all non-pruned stems to waste after final pruning
25-30	Clearfell		

Framing

Year	Operation	Stems per ha	Details
0	Establish	1000 - 1250	Improved genetics
8-10	Thin to waste	500-600	Thin all non-dominant stems to waste when mean crop height 14m
25-30	Clearfell		

Forest Health

Forest health surveys will be undertaken during routine forest visits. The Manager will complete its own health assessments.



Inventory, Mapping and Forest Records

Forest growth and development is monitored through regular forest inventory. Forest inventories providing stand information are required at different times and for different reasons throughout the life of a rotation. The Manager undertakes audits of all inventory to ensure consistency in approach and accuracy.

The following inventory is applied to Wairarapa Estate:

- Pre-assessment
- Quality Control
- Mid-rotation Inventory
- Pre-harvest Inventory

Pre-assessment

Pre-assessment is the collection of parameters prior to a tending operation to help calculate contract rates for tending, and to take a final check on the timing of the operation.

Sampling intensity is low with a minimum of five plots per stand, and data is collected from six to ten trees per plot. Data collected is then used to calculate a labour-day target and hence a contract rate per hectare. Contract rates are often set by tender or negotiation, reducing the need to pre-assess every block.

Pre-assessment is completed on the forests prior to tending operations commencing.

Quality Control

Quality control is carried out during and after a tending operation. The aims of the quality control are to:

- Collect sufficient data to monitor a contractor's performance and correct performance if necessary
- Collect quantitative data to provide reliable estimates of the crop
- Provide data as input for growth modelling

Sampling intensity is a minimum of five plots per stand, or one plot per 2 hectares, with every fourth plot being a full measurement plot. This provides the data for the current crop status and future growth modelling.

Data is summarised by Forest/Compartment/Stand prior to being entered into GeoMaster where it is retained as a permanent record. The records can then be directly accessed for annual reports and valuations and stand growth simulation modelling.

Quality control plots are completed at the stand level at the completion of each tending operation.

Mid-rotation Inventory

The principal aim for the mid-rotation inventory is to collect stand data for inputs into estate modelling and long-term harvest planning and marketing. The objective is to get accurate stand data summaries which will be used for crop typing, estate modelling and valuation.

This is a low intensity inventory, but with full log type cruising. This will enable summary to stand level and more accurate yield projections for the estate model. Mid-rotation inventory is scheduled for between 12 and 16 years of age.

Pre-harvest Inventory

The principal aim of pre-harvest inventory is to obtain estimates of recoverable volume by log grade. This information can then be used to develop marketing and harvesting strategies. Inventories will be undertaken when stands reach five years or less from harvesting. Sampling intensity is targeted to achieve 10% confidence limits on Basal Area on a stand-by-stand basis. Smaller stands may be aggregated into crop types to achieve this as in mid-rotation inventory.

Post-harvest Reconciliation

After harvest, reconciliation of data of the harvest area is undertaken to help improve records and to ensure harvesting has met the standards expected.



Mapping

Updating forest maps is required periodically as the forest changes. The work involves:

- Updating topographic detail
- Remapping forest stand boundaries from aerial photography
- Updating stand and forest attributes such as roads, landings, protected ecosystems and archaeological sites
- Defining legal boundaries

The data is kept and managed in the Manager's GIS system.

Stands are remapped from new aerial photographs around the age of four, when the trees are visible, to accurately determine boundaries. They are also remapped within 2 years of harvest to assist with harvest planning.

Forest Records

Forest records are essential to provide an historic perspective on the physical condition of each stand.

Forest records should provide the following information:

- Record of forest operations for each stand including a summary of quality control data
- · A forest map showing the location, stand boundaries and net stocked area of each stand
- Crop inventory results
- Yields achieved from each stand at production thinning or clearfell
- Costs incurred for each operation
- Protected ecosystems attributes
- Threatened species records
- Archaeological and Waahi Tapu sites and other potential features
- Chemicals used

The Manager maintains forest records in GeoMaster.



Harvesting

Currently, harvesting is scheduled by age class. However, as the estate grows, the Manager will attempt to smooth the harvest profile to ensure continuous harvesting with a similar volume harvested per annum.

Harvest Planning

All harvest planning and operations will follow the Manager's Environmental Standards.

Planning is essential to ensure roading infrastructure is developed in a timely manner and any resource consents and surveys are completed on time.

Harvest planning must consider:

- Slope determining what equipment can be used
- The Resource Management Act 1991, the Heritage NZ Pouhere Taonga Act 2014 and any other relevant legislation
- Safety how to ensure the operation is completed in a safe and legal manner
- Soil and water how to avoid, remedy or mitigate impacts on soil and water
- Ecosystems potential rare or threatened species and how to manage, managing in accordance with the New Zealand Forest Accord (1990) and the Code of Environmental Practice (ECoP)
- Possible sites of cultural, architectural, historical, ecological economic or religious significance to Tangata Whenua Tangata Whenua are to be consulted in regard to all planned harvest sites
- Financial outcomes plan the operation to ensure it meets current market demand and provides a return on investment
- Offsite impacts plan to minimise any adverse impacts on people or the environment outside of the forest

Harvesting Operations

All operations will be undertaken by harvest and transport contractors who have been selected for their quality of service and understanding of the **Manager's** Environmental Standards, associated SOPs, Health and Safety Management Systems (HSMS) and the Archaeological & Waahi Tapu Sites and Discovery SOP.

All operations will be supervised by the Manager, who has the right to stop the operation at any time if they feel the operation is having, or has the potential to have, an adverse impact on safety or the environment.

All operations will be regularly audited as per the Manager's health and safety and environmental systems.

Harvesting Methods

The Manager uses three basic criteria to ensure the right harvesting methods are employed:

- 1. Health & Safety the method is the most appropriate for the topography and nature of land so that the potential for injury is minimised
- 2. Environment the method creates the least impact on the environment
- 3. Financial the method is the most cost effective for the area taking safety and environmental considerations into account

The Manager is committed to adopting harvesting techniques and technology which minimise the impact on the environment and reduce the risk of accidents and injuries. To meet these objectives, Wairarapa Estate has been divided into four terrain types and the appropriate machinery configurations are used on each type.

Chain of Custody

All harvest loads leaving the estate will be accompanied by a docket or dockets stating crew, grade, forest location, weight and transport operator. Trucks will be randomly checked to ensure dockets are always present. The logs will be branded with the owners name.

Subject to attaining FSC certification, the Manager would ensure all dockets have the FSC certification number on them and may brand the logs with the FSC® logo.



Protection and Maintenance

The Manager will maintain roads, tracks, fences and water systems. The Manager will ensure pest and disease control, fire protection and management of protected areas occurs at all times.

Pest management within the estate is subject to statutory obligations under the Regional Pest Management Strategy administered by the Regional Council. The strategy applies to both pest plants and animals, categorising them in terms of management objectives. The categories, objectives and landowner obligations are summarised below for each Regional Pest Management Strategy Plan.

Weeds

The overall objective in managing weeds is to:

- Meet statutory obligations under the Regional Pest Management Strategy
- Reduce direct impacts on both plantations and indigenous biodiversity values
- Ensure impacts on neighbouring properties are promptly dealt with
- · Reduce the abundance and distribution of these species within the forest estate

The major species within Wairarapa Estate are various grasses, gorse, broom, blackberry and wilding conifers.

Competition from colonising weeds will limit tree growth in their first few years after establishment. Control of these weeds involves chemical application which will occur prior to planting and may occur post planting.

Gorse and broom threaten indigenous biodiversity in open communities where they can smother native species, however they can also act as a nurse crop in some areas for native regeneration. Blackberry can displace native species by outcompeting and smothering them.

A list of pest weeds appears in the appendix of the **Manager's** Environmental Standards and in the Appendix of this Plan.

All weeds will be controlled within Wairarapa Estate as follows.

(A) Total Control

Any weeds discovered under this category must be reported to the regional council. The council is responsible for controlling these weeds.

(B) Regional Surveillance

Weeds under this category should be reported to the council so the council can monitor them.

(C) Containment

Weeds under this category should be contained and removed if possible

(D) Site-led

These plants will be cleared by the Manager if within 50m of a neighbour.

Disease

Diseases, which can affect the forest trees and adjacent native vegetation, are monitored throughout the year. Most diseases can cause little damage and do not require control. The exception is *Dothistroma spp.*, a fungus which attacks pine needles and is associated with wet, warm conditions.

Dothistroma is the most commonly occurring fungal disorder within New Zealand's pine plantation. This fungus is controlled using an aerially applied copper-based fungicide spray, but only when the infection reaches a critical level. Dothistroma infection can also be controlled via silviculture by timely thinning and pruning operations, which increases air movement and lowers humidity levels.

There has been no need for Dothistroma control within the Wairarapa Estate forests.

No control is currently completed on other fungal disorders.

Any unusual mortality or colouring discovered within Wairarapa Estate will be reported to the Ministry for Primary Industries.



Chemical Control

All chemical applications are managed in accordance with the Manager's Environmental Standards, the Manager's Pesticide Application SOP and Chemical Use Policy, the New Zealand Standard for Agrichemical Application, Hazardous Substances and New Organisms Act 1996 regulations, and the obligations conferred by FSC to manage and minimise the use of chemicals; including use of alternatives where available.

As part of the FSC commitment:

- All chemical usage is tracked by active ingredient and application area to enable reporting and monitoring of trends
- The Manager has applied to be a participant in the Forest Owners environmental group which is undertaking research into chemical reduction, efficacy and safety issues related FSC regulations.
- The FSC Pesticides Policy (FSC-POL-30-001 V3-0 EN) is to be followed.
- All unwanted chemical can be booked for pick up via the Agrecovery website
- All empty containers will be disposed of at a registered recycling site

Pest Animals

Forests provide habitat for unwanted pest animals, and in most cases a refuge from which such pests can spread. Animal pests can substantially reduce the productivity of the forest.

The Manager will attempt to identify all pests present and manage them within the relevant regional council pest management strategies. The Manager will work to control or eradicate such species in accordance with these plans, to prevent spread and nuisance to neighbouring properties.

The most cost-effective long-term control is often achieved with the co-operation of neighbours, regional authorities and pest control agencies. The Manager will keep all these stakeholders informed of pest management operations.

Fire Prevention

The Manager complies with Fire and Emergency New Zealand (FENZ) and Wairarapa **Estate's fire plan which is** reviewed by the Manager each year before October.

The threat of fire is minimised by:

- Having an effective fire plan which encompasses prevention, detection and control procedures
- Active prevention measures which include restrictions on access, fire prevention signage, publicity when fire danger is high and access to water sources
- Effective detection systems which includes good communication systems, mapping, and fire plan alert procedures
- A close link with the relevant fire authorities, and an understanding of equipment and trained manpower availability
- Good forest management which recognises the influence of terrain, the road network and accessibility on fire prevention and control measures
- Suitable internal access systems of roads and tracks, and maintenance of fire breaks as the need arises

The legal responsibility for fighting forest fires lies with FENZ. In the event of a fire which starts within the forest, FENZ is responsible for attending and providing the resources to extinguish the fire.

The Manager maintains a close liaison with FENZ in terms of developing the "fire plan" and the maintenance of good communication relative to potential risks and fire danger ratings.

All neighbours are contacted prior to the fire season to check the Manager's records of contact numbers and other details are correct.



Rare, Threatened and Endangered Species

The Manager is committed to managing Wairarapa Estate to maintain a diversity of both indigenous flora and fauna species. Of particular importance are rare, threatened and endangered species living within the estate.

The Manager is undertaking a review to identify all rare, threatened and endangered species either confirmed or suspected to be present in the estate. Management Plans are progressively developed for all species confirmed to be present, focusing initially on those areas where harvesting is imminent.

Any permanent habitat for rare, threatened and endangered species is recorded in the GIS mapping layer as ecological restrictions, and taken into account during planning of operations, to ensure compliance with the Management Plans.



Monitoring

Every year, the Manager and Wairarapa Estate Limited will discuss the plan and associated annual plans. A review of measures taken to meet objectives will be undertaken at this time and outcomes recorded.

All monitoring will follow the Manager's monitoring plans and SOPs as per the Environmental Standards. Unless commercially sensitive, all monitoring results will be made available to public on request.

The monitoring program is designed to understand the impact of forest activities on the environment and the impact of the environment on the Manager's ability to grow trees. This leads to the development of strategies to ensure the Manager continues to manage its activities in a sustainable way.

Discussion will be held on the following:

Health and Safety

All contractors and staff will be audited as per the Manager's Health & Safety Management System (HSMS). All near misses, incidents and property damage will be recorded. The Manager will run a random drug and alcohol sampling program.

Environmental

The Manager will ensure all monitoring occurs as specified in its Environmental Standards. In particular, the Stream Health Monitoring and Assessment Kit (SHMAK) testing and Wildlands Ltd reporting on reserves.

Operations

The Manager conducts internal environmental audits to confirm operations have been carried out according to prescriptions, the Manager's Environmental Standards and regulatory requirements. Corrective actions are identified and rectified.

Regional Councils will also conduct resource consent compliance monitoring of operations undertaken under resource consents or permitted activity rules.

Forest Growth

Forest growth will be monitored through a combination of permanent sample plots and regular inventory.

Financial

The Manager will monitor budget versus expenditure quarterly and report variances to Wairarapa Estate Limited within the quarterly report and at quarterly meetings.

Annual reports will be provided, and periodic review meeting will be held when requested.

Forest Health

The Manager will undertake ongoing health monitoring survey to identify any health issues in the growing stands such as disease, pest damage or nutrient deficiencies.

The most common disease affecting radiata pine is a fungal disease *Dothistroma spp.* which causes needle cast in radiata pine and can severely slow tree growth. Dothistroma is controlled using copper-based products (cuprous oxide) similar to those used to control disease in home vegetable gardens.

Routine forest health inspections will identify any significant outbreaks of Dothistroma and be used to develop the annual spray programme. Significant amounts of research have been carried out to ensure the lowest possible effective level of fungicide is used to control this disease. Even though risk is low, application is planned to ensure that drift is minimised, and records are kept and audited to ensure that practices can be improved.

Stakeholders

Consultation will occur with stakeholders as per the **Manager's** Environmental Standards and this Forest Management Plan. Feedback from stakeholders will be sought and monitored. This includes actions undertaken to resolve disputes and issues, monitoring of externally generated complaints and client satisfaction surveys.

Consultation will occur with stakeholders during resource consent applications, annual and periodic meetings, contributions to council processes and interactions with forest recreational users and iwi.



Planning

This Forest Management Plan pertains to the management of Wairarapa Estate and will be adhered to for the next 5 years. Any deviation from this Plan will be justified only on the basis the changes do not adversely affect the environment. Any changes, which are contrary to the policies contained in this Plan require a full review of the Plan.

The next review date for this Plan is: March 2022.

The review will include review of planned monitoring, reserve areas and their protection, stakeholder engagement and financial performance.

The Forest Management Plan is used for both medium- and long-term planning.

For short-term operational and budgetary control planning, operations plans are prepared on an annual or as necessary basis. These plans are prepared annually and in accordance with this management plan. Operations plans and associated budgets are subject to approval by Wairarapa Estate Limited at the beginning of each financial year.



Appendices

- 1. Pest Plants of the Wellington Region
- 2. Relevant Regulations, Standards and Guidelines
- 3. Maps and Management Plans Botanised High Conservation Value Areas
- 4. Maps High Conservation Value Areas
- 5. Maps Stream Health Monitoring and Assessment (SHMAK) Sites
- 6. Maps Erosion Susceptibility Classification (ESC)



Appendix 1: Pest Plants of the Wellington Region

African fountain grass Pennisetum setaceum Key Native Ecosystem Programme Alligator weed Alternanthera philoxeroides Exclusion Apple of Sodom Solanum linnaeanum Key Native Ecosystem Programme Artemisia Artemisia spp. Key Native Ecosystem Programme Artillery plant Galeobdolon luteum Key Native Ecosystem Programme Arum lily Zantedeschia aethiopica Key Native Ecosystem Programme Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Australian sedge Carex longebraciata Key Native Ecosystem Programme Banana passionfruit Passiflora mixta, P. mollissima, P. tripartita Site-led Hutt City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Froom Cytisus scoparius Key Native Ecosystem Programme Key Native Ecosystem Programme Froom Cytisus scoparius Key Native Ecosystem Programme Froom Cytisus scoparius Key Native Ecosystem Programme Froom Cytisus scoparius Key Native Ecosystem Programme Froom Reynamine Froormame	Plants	Scientific Name	Primary Programme
African fountain grass Pennisetum setaceum Alternanthera philoxeroides Exclusion Apple of Sodom Aple of Sodom Solanum linnaeanum Key Native Ecosystem Programme Key Native Ecosystem Programme Key Native Ecosystem Programme Artemisia Artemisia Artemisia spp. Artillery plant Galeobdolon luteum Key Native Ecosystem Programme Arum iliy Zantedeschia aethiopica Key Native Ecosystem Programme Arum iliy Zantedeschia aethiopica Key Native Ecosystem Programme Arum iliy Zantedeschia aethiopica Key Native Ecosystem Programme Australian sedge Carex longebraciata Key Native Ecosystem Programme Australian sedge Carex longebraciata Key Native Ecosystem Programme Banana passionfruit Passiffora mixta, P. mollissima, P. tripartita Site-led Hutt City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiffora caerulea Sustained control Bomarea Bomarea Bomarea caidasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytius scopanius Key Native Ecosystem Programme Broom Cytius scopanius Key Native Ecosystem Programme Buddleia Buddleia Buddleia Buddleia Buddleia Buddleia davidii Key Native Ecosystem Programme Californian burtush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape tulp Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cape tulp Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Key Native Ecosystem Programme Chilean needle grass Asparagus scandens Key Native Ecosystem Programme Chilean needle grass Asparagus scandens Key Na	African club moss	Selaginella kraussiana	Key Native Ecosystem Programme
Alternanthera philoxeroides Exclusion Apple of Sodom Solanum linnaeanum Key Native Ecosystem Programme Artemisia Artemisia spp. Key Native Ecosystem Programme Artillery plant Galeobdolon luteum Key Native Ecosystem Programme Artillery plant Galeobdolon luteum Key Native Ecosystem Programme Arum Iliy Zantedeschia aethiopica Key Native Ecosystem Programme Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Australian sedge Carex longebraciata Key Native Ecosystem Programme Australian sedge Carex longebraciata Key Native Ecosystem Programme Banana passionfruit Passiffora mixta, P. mollissima, P. tripartita Site-led Hutt City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiffora caerulea Sustained control Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Progr	African feather grass	Pennisetum macrourum	Key Native Ecosystem Programme
Apple of Sodom Solanum linnaeanum Key Native Ecosystem Programme Artemisia Artemisia Artemisia spp. Key Native Ecosystem Programme Artillery plant Galeobdolon luteum Key Native Ecosystem Programme Arum iliy Zantedeschia aethiopica Key Native Ecosystem Programme Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Banana passionifuit Passiflora mixta, P. mollissima, P. tripartita Bite-led Hutt City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Broam Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Californian arrowhead Sagiltaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Chieban needle grass Nassella neesiana Exclusion Chineae pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chieban gasparagus Asparagus Scandens Key Native Ecosystem Programme Climbing asparagus Asparagus Scandens Key Native Ecosystem Programme Climbing asparagus Cotoneaster Cotoneaster Cotoneaster Cotoneaster Coto	African fountain grass	Pennisetum setaceum	Key Native Ecosystem Programme
Artemisia Artemisia spp. Key Native Ecosystem Programme Artillery plant Galeobdolon luteum Key Native Ecosystem Programme Artum Iliy Zantedeschia aethiopica Key Native Ecosystem Programme Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Australian sedge Carex longebraciata Key Native Ecosystem Programme Banana passionfruit Passiflora mixta, P. mollissima, P. tripartita Site-led Hutt City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Sustained control Chrysanthemoides monilifera Sustained control Chrysanthemoides Mey Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape ivy Seneci	Alligator weed	Alternanthera philoxeroides	Exclusion
Artillery plant Galeobdolon luteum Key Native Ecosystem Programme Arum Iliy Zantedeschia aethiopica Key Native Ecosystem Programme Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Australian sedge Garek longebraciata Key Native Ecosystem Programme Banana passionfruit Passiflora mixta, P. mollissima, P. tripartita Site-led Hut City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Asanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Boneseed Chrysanthemoides monilifera Sustained control Chrysanthemoides Chrysanthemoides Rey Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleia Buddleia davidii Key Native Ecosystem Programme Californian butrush Schoenoplectus californicus Key Native Ecosystem Programme Californian butrush Schoenoplectus californicus Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape Linding Alexander Chrocolate vine Akebia quinata Key Native Ecosystem Programme Chicolate vine Akebia quinata Key Native Ecosystem Programme Chicolate vine Akebia quinata Key Nat	Apple of Sodom	Solanum linnaeanum	Key Native Ecosystem Programme
Arum lily Zantedeschia aethiopica Key Native Ecosystem Programme Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Australian sedge Care Iongebraciata Key Native Ecosystem Programme Banana passionfruit Passiflora mixta, P. mollissima, P. tripartita Site-led Hutt City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Bomarea Bomarea Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broseed Chrysanthemoides monilifera Sustained control Key Native Ecosystem Programme Brush wattle Paraserianthes Iophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chicolate vine Akebia quinata Key Native Ecosystem Programme Chicolate vine Akebia quinata Key Native Ecosystem Programme Cilimbing aparagus Asparagus scandens Key Native Ecosystem Programme Cilimbing aparagus Asparagus scandens Key Native Ecosystem Programme Cilimbing apindleberry Celastrus orbiculatus Key Native Ecosystem Programme Cilimbing spindleberry Celastrus orbiculatus Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme	Artemisia	Artemisia spp.	Key Native Ecosystem Programme
Asiatic knotweed Reynoutria japonica Key Native Ecosystem Programme Australian sedge Carex longebraciata Key Native Ecosystem Programme Banana passionfruit Passiflora mixta, P. mollissima, P. tripartita Site-led Hutt City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Key Native Ecosystem Programme Bonaseed Chrysanthemoides monilifera Sustained control Cyclium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes Iophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chicolate vine Akebia quinata Key Native Ecosystem Programme Chicilian needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Cilimbing asparagus Asparagus scandens Key Native Ecosystem Programme Cilimbing asparagus Asparagus scandens Key Native Ecosystem Programme Key Native Ecosystem Programme Cilimbing asparagus Asparagus scandens Key Native Ecosystem Programme Key Native Ecosystem Programme Cilimbing spindleberry Celastrus orbiculatus Sustained control Key Native Ecosystem P	Artillery plant	Galeobdolon luteum	Key Native Ecosystem Programme
Australian sedge Carex longebraciata Key Native Ecosystem Programme Banana passionfruit Passiflora mixta, P. mollissima, P. tripartita Site-led Hutt City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Bush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Cathedral bells Cobaea scandens Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Crack willow Salix fragilis	Arum lily	Zantedeschia aethiopica	Key Native Ecosystem Programme
Banana passionfruit Passiflora mixta, P. mollissima, P. tripartita Site-led Hutt City Council (HCC) Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Climbing asparagus Climbing asparagus Climbing asparagus Climbing asparagus Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Berberis darwinii Key Native Ecosystem Programme Crack willow Berberis darwinii Key Native Ecosystem Programme	Asiatic knotweed	Reynoutria japonica	Key Native Ecosystem Programme
Barberry Berberis glaucocarpa Key Native Ecosystem Programme Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes Iophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape lip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Australian sedge	Carex longebraciata	Key Native Ecosystem Programme
Bathurst bur Xanthium spinosum Key Native Ecosystem Programme Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape lulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Crack willow Berberry Berbis darwinii Key Native Ecosystem Programme	Banana passionfruit	Passiflora mixta, P. mollissima, P. tripartita	Site-led Hutt City Council (HCC)
Blackberry Rubus spp. barbed cultivars Key Native Ecosystem Programme Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Crack willow Berberry Berbis darwinii Key Native Ecosystem Programme	Barberry	Berberis glaucocarpa	Key Native Ecosystem Programme
Blue morning glory Ipomoea indica Key Native Ecosystem Programme Blue passionflower Passiflora caerulea Sustained control Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes Iophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Califomian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme	Bathurst bur	Xanthium spinosum	Key Native Ecosystem Programme
Blue passionflower Bomarea Bomarea aldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Buddleja davidii Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Key Native Ecosystem Programme Cimbing spindleberry Celastrus orbiculatus Sustained control Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme	Blackberry	Rubus spp. barbed cultivars	Key Native Ecosystem Programme
Bomarea Bomarea caldasii, B. multiflora Key Native Ecosystem Programme Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme	Blue morning glory	Ipomoea indica	Key Native Ecosystem Programme
Boneseed Chrysanthemoides monilifera Sustained control Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme	Blue passionflower	Passiflora caerulea	Sustained control
Boxthorn Lycium ferocissimum Key Native Ecosystem Programme Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Crack willow Berberis darwinii Key Native Ecosystem Programme	Bomarea	Bomarea caldasii, B. multiflora	Key Native Ecosystem Programme
Broom Cytisus scoparius Key Native Ecosystem Programme Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Cimbing spindleberry Celastrus orbiculatus Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberisi darwinii Key Native Ecosystem Programme	Boneseed	Chrysanthemoides monilifera	Sustained control
Brush wattle Paraserianthes lophantha Key Native Ecosystem Programme Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Cimbing spindleberry Celastrus orbiculatus Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Boxthorn	Lycium ferocissimum	Key Native Ecosystem Programme
Buddleia Buddleja davidii Key Native Ecosystem Programme Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Broom	Cytisus scoparius	Key Native Ecosystem Programme
Californian arrowhead Sagittaria montevidensis Key Native Ecosystem Programme Californian bulrush Schoenoplectus californicus Key Native Ecosystem Programme Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Key Native Ecosystem Programme Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme	Brush wattle	Paraserianthes lophantha	Key Native Ecosystem Programme
Californian bulrush Cape honey flower Melianthus major Key Native Ecosystem Programme Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Key Native Ecosystem Programme	Buddleia	Buddleja davidii	Key Native Ecosystem Programme
Cape honey flower Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme	Californian arrowhead	Sagittaria montevidensis	Key Native Ecosystem Programme
Cape ivy Senecio angulatus Key Native Ecosystem Programme Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Californian bulrush	Schoenoplectus californicus	Key Native Ecosystem Programme
Cape tulip Moraea flaccida (syn. Homeria collina) Key Native Ecosystem Programme Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Crack willow Berberis darwinii Key Native Ecosystem Programme	Cape honey flower	Melianthus major	Key Native Ecosystem Programme
Cathedral bells Cobaea scandens Site-led HCC Chilean flame creeper Tropaeolum speciosum Key Native Ecosystem Programme Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Crack willow Salix fragilis Key Native Ecosystem Programme	Cape ivy	Senecio angulatus	Key Native Ecosystem Programme
Chilean flame creeper Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Key Native Ecosystem Programme	Cape tulip	Moraea flaccida (syn. Homeria collina)	Key Native Ecosystem Programme
Chilean needle grass Nassella neesiana Exclusion Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Cathedral bells	Cobaea scandens	Site-led HCC
Chinese pennisetum Pennisetum alopecuroides Key Native Ecosystem Programme Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Chilean flame creeper	Tropaeolum speciosum	Key Native Ecosystem Programme
Chocolate vine Akebia quinata Key Native Ecosystem Programme Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Chilean needle grass	Nassella neesiana	Exclusion
Climbing asparagus Asparagus scandens Key Native Ecosystem Programme Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Chinese pennisetum	Pennisetum alopecuroides	Key Native Ecosystem Programme
Climbing dock Rumex sagittatus Key Native Ecosystem Programme Climbing spindleberry Celastrus orbiculatus Sustained control Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Chocolate vine	Akebia quinata	Key Native Ecosystem Programme
Climbing spindleberry Celastrus orbiculatus Sustained control Key Native Ecosystem Programme Crack willow Salix fragilis Berberis darwinii Sustained control Key Native Ecosystem Programme Key Native Ecosystem Programme Key Native Ecosystem Programme	Climbing asparagus	Asparagus scandens	Key Native Ecosystem Programme
Cotoneaster Cotoneaster franchetii, C. horizontalis Key Native Ecosystem Programme Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Climbing dock	Rumex sagittatus	Key Native Ecosystem Programme
Crack willow Salix fragilis Key Native Ecosystem Programme Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Climbing spindleberry	Celastrus orbiculatus	Sustained control
Darwin's barberry Berberis darwinii Key Native Ecosystem Programme	Cotoneaster	Cotoneaster franchetii, C. horizontalis	Key Native Ecosystem Programme
	Crack willow	Salix fragilis	Key Native Ecosystem Programme
Delta arrowhead Sagittaria platyphylla Key Native Ecosystem Programme	Darwin's barberry	Berberis darwinii	Key Native Ecosystem Programme
	Delta arrowhead	Sagittaria platyphylla	Key Native Ecosystem Programme



Didymo	Didymosphenia geminata	Key Native Ecosystem Programme
Eelgrass	Vallisneria spiralis, V. gigantea	Sustained control
Elaeagnus	Elaeagnus x reflexa	Key Native Ecosystem Programme
Evergreen buckthorn	Rhamnus alaternus	Key Native Ecosystem Programme
German ivy	Senecio mikanioides	Key Native Ecosystem Programme
Giant knotweed	Reynoutria sachalinensis and hybrids	Key Native Ecosystem Programme
Gorse	Ulex europaeus	Key Native Ecosystem Programme
Great bindweed	Calystegia silvatica	Key Native Ecosystem Programme
Gunnera	Gunnera tinctoria	Key Native Ecosystem Programme
Hawaiian arrowhead	Sagittaria sagittifolia	Key Native Ecosystem Programme
Hawthorn	Crataegus monogyna	Key Native Ecosystem Programme
Hemlock	Conium maculatum	Key Native Ecosystem Programme
Himalayan honeysuckle	Leycesteria formosa	Key Native Ecosystem Programme
Hornwort	Ceratophyllum demersum	Key Native Ecosystem Programme
Houttuynia	Houttuynia cordata	Key Native Ecosystem Programme
Hydrilla	Hydrilla verticillata	Key Native Ecosystem Programme
Japanese honeysuckle	Lonicera japonica	Key Native Ecosystem Programme
Japanese spindletree	Euonymus japonicus	Key Native Ecosystem Programme
Johnson grass	Sorghum halepense	Key Native Ecosystem Programme
Lagarosiphon	Lagarosiphon major	Key Native Ecosystem Programme
Madeira vine	Anredera cordifolia	Key Native Ecosystem Programme
Manchurian wild rice	Zizania latifolia	Key Native Ecosystem Programme
Marram grass	Ammophila arenaria	Key Native Ecosystem Programme
Mexican daisy	Erigeron karvinskianus	Key Native Ecosystem Programme
Mile-a-minute	Dipogon lignosus	Key Native Ecosystem Programme
Mist flower	Ageratina riparia	Key Native Ecosystem Programme
Monkey apple	Acmena smithii	Key Native Ecosystem Programme
Montbretia	Crocosmia x crocosmiflora	Key Native Ecosystem Programme
Moth plant	Araujia hortorum	Eradication
Nassella tussock	Nassella trichotoma	Exclusion
Nasturtium	Nasturtium officinalis	Key Native Ecosystem Programme
Nodding thistle	Carduus nutans	Key Native Ecosystem Programme
Noogoora bur	Xanthium occidentale	Key Native Ecosystem Programme
Old man's beard	Clematis vitalba	Site-led HCC
Pampas grass	Cortaderia jubata, C. selloana	Key Native Ecosystem Programme
Parrot's feather	Myriophyllum aquaticum	Key Native Ecosystem Programme
Perennial nettle	Urtica dioica (subspp.)	Key Native Ecosystem Programme
Periwinkle	Vinca major	Key Native Ecosystem Programme
Phragmites	Phragmites australis	Key Native Ecosystem Programme
Plectranthus	Plectranthus ciliatus	Key Native Ecosystem Programme
		, Loody old in rogidinino
Polypodium (common polypody)	Polypodium vulgare	Key Native Ecosystem Programme



Purple ragwort	Senecio glastifolius	Key Native Ecosystem Programme
Pussy willow	S. cinerea	Key Native Ecosystem Programme
Pyp grass	Ehrharta villosa	Key Native Ecosystem Programme
Ragwort	Senecio jacobaea	Key Native Ecosystem Programme
Saffron thistle	Carthamus lanatus	Key Native Ecosystem Programme
Salvinia	Salvinia molesta	Key Native Ecosystem Programme
Senegal tea	Gymnocoronis spilanthoides	Eradication
Silver poplar	Populus alba	Key Native Ecosystem Programme
Smilax	Asparagus asparagoides	Key Native Ecosystem Programme
Spanish heath	Erica lusitanica	Key Native Ecosystem Programme
Spartina	Spartina anglica, S. alterniflora	Eradication
Stinking iris	Iris foetidissima	Key Native Ecosystem Programme
Sweet pea shrub	Polygala myrtifolia	Key Native Ecosystem Programme
Sycamore	Acer pseudoplatanus	Key Native Ecosystem Programme
Tradescantia	Tradescantia fluminensis	Key Native Ecosystem Programme
Tuber ladder fern	Nephrolepis cordifolia	Key Native Ecosystem Programme
Variegated thistle	Silybum marianum	Key Native Ecosystem Programme
Velvet groundsel	Senecio petasitis	Key Native Ecosystem Programme
Velvetleaf	Abutilon theophrasti	Eradication
Water hyacinth	Eichhornia crassipes	Key Native Ecosystem Programme
White bryony	Bryonia cretica subsp. dioica	Key Native Ecosystem Programme
White edged nightshade	Solanum marginatum	Key Native Ecosystem Programme
Wild ginger	Hedychium, gardnerianum, H. flavescens	Key Native Ecosystem Programme
Wild onion	Allium vineale	Key Native Ecosystem Programme
Wilding conifers	Pinus spp., Macrocarpa spp.	Progressive containment
Woolly nightshade	Solanum mauritianum	Eradication



Appendix 2: Relevant Regulations, Standards and Guidelines

A.	NATIONAL LEGISLATION
	Accident Compensation Act 2001
	Anti-Money Laundering and Countering Financing of Terrorism Act 2009
	Biosecurity Act 1993
	Climate Change (Forestry Sector) Regulations 2008
	Climate Change Response Act 2002
	Code of Practice for the Management of Agrichemicals 2004 (NZS8409:2004)
	Commerce Act 1986
	Companies Act 1993
	Conservation Act 1987
	Contract and Commercial Law Act 2017
	Cooperative Companies Act 1996
	Crown Forest Assets Act 1989
	Crown Minerals Act 1991
	Ecological Regions and Districts of New Zealand
	Employment Relations Act 2000
	Fair Trading Act 1986
	Fencing Act 1978
	Fire and Emergency New Zealand Act 2017
	Forest Disease Control Regulations 1967
	Forest Produce Import & Export Regulations 1989
	Forestry Encouragement Act 1962
	Forestry Encouragement Loans Regulations 1967
	Forestry Rights Registration Act 1983
	Forests Act 1949
	Forests (Regulation of Log Traders and Forestry Advisers) Amendment Act 2020
	Goods and Services Tax Act 1985
	Hazardous Substances & New Organisms Act 1996
	Health & Safety at Work Act 2015
	Heritage New Zealand Pouhere Taonga Act 2014
	Holidays Act 2003
	Income Tax Act 2007
	Land Act 1948
	Land Transfer Act 1952
	Land Transport Act 1998
	Local Government Act 2002
	Machinery Act 1950
	Maori Reserved Land Act 1955
	Marine and Coastal Area (Takutai Moana) Act 2011
	Minimum Wage Act 1983



Misuse of Drugs Act 1975		
National Environmental Standards for Plantation Forestry		
Native Plants Protection Act 1934		
New Zealand Environmental Code of Practice for Plantation Forestry 2007		
New Zealand Forest Code of Practice June 1993		
New Zealand Threat Classification System 2005		
Overseas Investment Act 2005		
Personal Property Securities Act 1999		
Principles for Commercial Plantation Forest Management in New Zealand 1995		
Privacy Act 2020		
Public Works Act 1981		
Resource Management Act 1991		
Safety and Health in Forestry Operations: Code of Practice and Best Practice Guidelines		
Soil Conservation and Rivers Control Act 1941		
The New Zealand Forest Accord 1991		
Trade Marks Act 2002		
Treaty of Waitangi Act 1975		
Trespass Act 1980		
Walking Access Act 2008		
Wild Animal Control Act 1977		
Wildlife Act 1953		
Legal Rights to Harvest:		
Land tenure and management rights		
Concession licenses		
Management and harvest planning		
Trade and Transport		
 Classification of species, quantities, qualities Trade and transport 		
Offshore trading and transfer pricing		
Third Party Rights		
Customary rights		
Free prior and informed consent (FPIC)		
Rights of indigenous peoples		
Timber Harvesting Activities		
Timber harvesting regulations		
Protected sites and species		
Environmental requirements - Health and acfatr.		
Health and safetyLegal employment		
Taxes and Fees		
Payment of royalties and harvesting fees		
Value added and sales taxes		
Income and profit taxes		
Custom Regulations		
Biosecurity Act 1993		



	Customs and Excise Act 2018	
	Forests Act 1949 The New Zealand Forest Accord 1991	
	CITES	
	Convention on the International Trade in Endangered Species (CITES)	
	Other	
	Not applicable at this stage. All relevant legislation has been stated	
В.	REGULATIONS PERTINENT TO FORESTRY RELATED TO AND EMERGING FROM NATIONAL LEGISLATION AND OTHER LEGISLATIVE INSTITUTIONS:	
	Code of Practice for the Management of Agrichemicals 2004 (NZS8409:2004)	
	Ecological Regions and Districts of New Zealand	
	Fire and Emergency New Zealand Act 2017	
	Fire and Emergency New Zealand Regulations	
	Forestry Encouragement Loans Regulations 1967	
	Forestry Rights Registration Act 1983	
	Forests Act 1949	
	New Zealand Environmental Code of Practice for Plantation Forestry 2007	
	New Zealand Forest Code of Practice June 1993	
	New Zealand Threat Classification system 2005	
	Principles for Commercial Plantation Forest Management in New Zealand 1995	
	Resource Management Act 1991	
	The New Zealand Forest Accord 1991	



C.	INTERNATIONAL AGREEMENTS PERTINENT TO FORESTRY
	Convention on Biological Diversity
	Convention on the International Trade in Endangered Species (CITES)
	ICOMOS New Zealand Charter 1993
	ITTA
	IUCN Red List of threatened species
	Kyoto Protocol
	International Labour Organisation (ILO) conventions: 29 Forced Labour Convention 1930 87 Freedom of Association and Protection of the Right to Organise Conventions 1948 97 Migration for Employment (Revised) Convention 1949 98 Right to Organise and Collective Bargaining Convention 1949 100 Equal Remuneration Convention 1951 105 Abolition of Forced Labour Convention 1957 111 Discrimination (Occupation and Employment) Convention 1958 131 Minimum Wage Fixing Convention 1970 138 Minimum Age Convention 1973 141 Rural Workers' Organizations Convention 1975 142 Human Resources Development Convention 1975 143 Migrant Workers (Supplementary Provisions) Convention 1975 155 Occupational Safety and Health Convention 1981 169 Indigenous and Tribal Peoples Convention 1989 182 Worst Forms of Child Labour Convention 1999 ILO Code of Practice on Safety and Health in Forestry Work (ILO 1998) Recommendation 135 Minimum Wage Fixing Recommendation 1970
	 ILO Declaration on Fundamental Principles and Rights at Work 1998 and its follow-up ILO member states are expected to promote and realize these principles, even if they have not ratified the Conventions
	 The ILO Code of Practice is not a legal instrument, but it provides authoritative guidance on forest work ILO Code of Practice is not a legal instrument, but it provides authoritative guidance on forest work
D.	LOCAL STANDARDS AND BEST OPERATING PRACTICES
	Code of Practice for the Management of Agrichemicals 2004 (NZS8409:2004)
	Ecological Regions and Districts of NZ
	National Environmental Standards for Plantation Forestry
	New Zealand Environmental Code of Practice for Plantation Forestry 2007
	New Zealand Forest Code of Practice June 1993
	New Zealand Threat Classification system 2005
	Principles for Commercial Plantation Forest Management in New Zealand 1995
	Safety and Health in Forestry Operations: Code of Practice and Best Practice Guidelines
	The New Zealand Forest Accord 1991



Appendix 3: Maps and Management Plans – High Conservation Value Forest Areas

HCVF areas are monitored annually and ecological surveys occur every two years.

Saline Springs in Glenburn Forest

The High Conservation Value Forest (HCVF) area is to be protected with a deer fence erected around the actual springs. There will be a gate, on which the Manager will place a lock.

The Manager will remove all exotic vegetation from within and around the fence.

Access to the Springs will be limited.

A photo point will be set up within the HCVF area to help with monitoring. Monitoring will occur annually. The HCVF area will be monitored for:

- Weed growth, which the Manager will remove
- Native growth of saline species




Appendix 4: Maps – High Conservation Value Areas




Beehive Creek High Conservation Value Map

1:7,000











Craigie Lea High Conservation Value Map

1:25,000











Driscoll Road High Conservation Value Map

1:7,000









Dunolly High Conservation Value Map

1:15,000











Erindale High Conservation Value Map

1:17,000











Flat Point High Conservation Value Map

1:6,000











Glenburn (North) High Conservation Value Map

1:26,000










Glenburn (South) High Conservation Value Map


1:27,000



Date: 24/02/2021



Wairarapa Estate Ltd




Hapua High Conservation Value Map

1:14,000




Date: 24/02/2021



Silver Range

Elsthorpe

Patangata




Hawkins High Conservation Value Map

1:8,000




Date: 24/02/2021



Carswell Awatoitoi

Tauweru River

Mangapakeha Mangapakeha Taipo



Kaiwhata Pines High Conservation Value Map

1:11,000




Date: 24/02/2021



Te Wharau


Pukeroro Mapapa Stream




Lands End High Conservation Value Map

1:6,000










Ngahape - WE High Conservation Value Map

1:9,000











Oldfields High Conservation Value Map

1:9,000












Pakowhai High Conservation Value Map

1:19,000











Pongaroa High Conservation Value Map

1:13,000












Putorino High Conservation Value Map

1:13,000









Riverina High Conservation Value Map

1:8,000




Date: 24/02/2021



Frasertown


Tutaekuri River




Roil High Conservation Value Map

1:8,000










Ruakokoputuna High Conservation Value Map

1:13,000











Tinui - WE High Conservation Value Map

1:14,000











Wai Ngaio High Conservation Value Map

1:9,000










Waipukurau High Conservation Value Map

1:4,000











Woodford Green High Conservation Value Map

1:7,000









Appendix 5: Maps – Stream Health Monitoring and Assessment (SHMAK) Sites




Beehive Creek SHMAK Map

1:7,000



Date: 22/02/2021



Pohangina Awahou North

Pohangina River

Awahou South


Wairarapa Estate Ltd

Cheltenham

Almadale

Colyton

Hiwinui



Craigie Lea SHMAK Map

1:25,000



Date: 22/02/2021



Honeycomb Rock Waihinga

Wairarapa Estate Ltd

Public Road - Unmetalled Contour


SHMAK Site

==== Forest Road

Rerewhakaaitu River Stocked

NonProductive

20m (Index)



Driscoll Road SHMAK Map

1:7,000



Date: 22/02/2021



Ngãumu Forest

Kaiwhata River

Te Wharau


Mapapa Stream

Rocky Hills Sanctuary Area

Pahaoa River

Maungarake Range


River Spur




Dunolly SHMAK Map

1:15,000










1:17,000



Date: 22/02/2021



Mataikona

Whakataki River

Whakataki


Castlepoint

Whareama River

Tinui River

Langdale


Whareama




Flat Point SHMAK Map

1:6,000







Glenburn (North) SHMAK Map

1:26,000




Date: 22/02/2021



Stocked

Wairarapa Estate Ltd



Glenburn (South) SHMAK Map


1:27,000



Date: 22/02/2021




Wairarapa Estate Ltd





Hapua SHMAK Map

1:14,000









Hawkins SHMAK Map

1:8,000



Date: 22/02/2021

Stocked


Landbank
NonProductive

Mangapakeha

Whareama


Mangapakeha Taipo

Carswell Awatoitoi



Wairarapa Estate Ltd

SHMAK Site




Kaiwhata Pines SHMAK Map

1:11,000



Date: 22/02/2021




Riversdale Beach

Ngāumu Forest

Westmere

Mapapa Stream

Te Wharau




Lands End SHMAK Map

1:6,000



Date: 22/02/2021




Stocked Landbank NonProductive

Contour

Wairarapa Estate Ltd

Ground Based



Ngahape - WE SHMAK Map

1:9,000



Date: 22/02/2021



Meters

Whakatahine River

Wainuiora River

Kaiwhata River

Pukeroro


Ngãumu Forest

Riversdale Beach

220

Te Wharau


Westmere




Oldfields SHMAK Map

1:9,000














1:19,000











Pongaroa SHMAK Map

1:13,000








Putorino SHMAK Map

1:13,000




Date: 22/02/2021



NonProductive

Wairarapa Estate Ltd
S SHMAK Site





Riverina SHMAK Map

1:8,000










Roil SHMAK Map

1:8,000



Date: 22/02/2021



The Blowhard Ridge

Longfellow Range Sandy Ridge

Otamauri

Glenross Range The Lizard Willowford Waikarikari

Stocked


Wairarapa Estate Ltd

Hoodoo Saddle

Sparrowhawk Range

Kaikomata Range Glapros

> Hollowback Ridge




Ruakokoputuna SHMAK Map

1:13,000



Date: 22/02/2021



Pukio Dry River

Ruakokoputuna River


Castle River

Tuhitarata

Aorangi Range

Huangarua River

Ruakokoputuna Makara River




Tinui - WE **SHMAK Map**

1:14,000



Date: 22/02/2021



Whareama


Blairlogie

NonProductive

20m (Index)

Wairarapa Estate Ltd

Public Road - Sealed




Wai Ngaio SHMAK Map

1:9,000



Date: 22/02/2021



Tinui River

Langdale


Waimimi

Mangapakeha Mangapakeha Taipo

Blairlogie Whareama

Whakataki


Castlepoint




Waipukurau SHMAK Map

1:4,000









Woodford Green SHMAK Map

1:7,000









Appendix 6: Maps – Erosion Susceptibility Classification (ESC)



